

Design and Evaluation of a Mobile Chat App for the Open Source Behavioral Health Intervention Platform MobileCoach

Tobias Kowatsch¹(✉), Dirk Volland², Iris Shih², Dominik Rügger², Florian Künzler²,
Filipe Barata², Andreas Filler^{1,3}, Dirk Büchter⁴, Björn Brogle⁴, Katrin Heldt⁴, Pauline
Gindrat⁵, Nathalie Farpour-Lambert⁶ & Dagmar l'Allemand⁴

¹ University of St. Gallen, Institute of Technology Management, St. Gallen, Switzerland
tobias.kowatsch@unisg.ch

² ETH Zurich, Department of Management, Technology and Economics, Zurich, Switzerland

³ University of Bamberg, Energy Efficient Systems Group, Bamberg, Germany

⁴ Children's Hospital of Eastern Switzerland, St. Gallen, Switzerland

⁵ Fondation SportSmile, Nyon, Switzerland

⁶ University Hospital of Geneva / University of Geneva, Department of Community Medicine,
Primary Care and Emergency, Geneva, Switzerland

Abstract. The open source platform MobileCoach (mobile-coach.eu) has been used for various behavioral health interventions in the public health context. However, so far, MobileCoach is limited to text message-based interactions. That is, participants use error-prone and laborious text-input fields and have to bear the SMS costs. Moreover, MobileCoach does not provide a dedicated chat channel for individual requests beyond the processing capabilities of its chatbot. Intervention designers are also limited to text-based self-report data. In this paper, we thus present a mobile chat app with pre-defined answer options, a dedicated chat channel for patients and health professionals and sensor data integration for the MobileCoach platform. Results of a pretest ($N=11$) and preliminary findings of a randomized controlled clinical trial ($N=14$) with young patients, who participate in an intervention for the treatment of obesity, are promising with respect to the utility of the chat app.

Keywords: Health intervention • Digital coaching • Chat-based interaction.

1 Introduction

Non communicable diseases (NCDs) such as heart diseases, asthma, obesity, diabetes or chronic kidney disease impose the greatest burden on global health [14]. According to WHO's NCD global monitoring framework, many of these diseases are consequences of adverse health behaviors, for example, harmful use of alcohol and tobacco or physical inactivity [15]. However, health personnel is strongly limited [2]. Consequently, scalable behavioral health interventions are required.

Innovative digital health interventions (DHIs) have not only the potential to improve the efficacy of preventive or therapeutic behavioral health interventions but also to reduce their costs [1]. With the goal to provide an open source platform that allows health professionals to design scalable, low-cost and evidence-based DHIs, MobileCoach (mobile-coach.eu) was developed [5] and evaluated [9].

However, it uses the short message service (SMS) for delivering behavioral health interventions, and thus comes with various shortcomings as outlined in the next section. We therefore present in this paper the first mobile chat app for the MobileCoach platform that addresses these shortcomings and thus, complements existing communication such as personal exchange, SMS-based, phone-based or video-based interactions.

The remainder of this paper is structured as follows. Next, we describe the design of the chat app. Then, the app's significance to research and practice is outlined. Finally, we present results from an empirical study with 11 obese children who assessed the new chat app as the first target group.

2 Design of the Chat App

Hands-on experience with several MobileCoach-based interventions [7-9] has revealed four major shortcomings related to its text messaging approach. First, participants have to bear the SMS costs which may be an entry barrier if the caregiver does not provide a monetary compensation. Second, participants are always requested to manually type in text to answer even Likert-scale type questions. These answers are then parsed by the MobileCoach, which is error-prone in case the answer does not perfectly fit to the question. Processing these answers is a time-consuming process for the caregiver, too. Third, participant-initiated requests usually require an individual answer from a caregiver instead of a scripted answer by a chatbot. A rule-based chatbot does therefore not always fit to the communication needs of the participants. Fourth, text-messaging is limited to self-report data, i.e. health professionals cannot use objective sensor data from a smartphone (e.g., accelerometer data used to measure physical activity) or sensor data from devices connected to that smartphone (e.g., Bluetooth-enabled blood glucose or peak flow meters) for the design of their DHIs.

Regarding these shortcomings and against the background of smartphone pervasiveness [4], the following requirements have been defined: (R1) The app must not rely on the short message service for communication purposes; (R2) the app must implement pre-defined answer sets for efficient and error-free chat interaction; (R3) the app must implement a chat channel for individual communication needs that complements the scalable chatbot channel; (R4) the app must be able to access sensor data from the smartphone or smartphone-connected devices.

By considering these four requirements, we built a first mock-up of a mobile app and evaluated it with six behavioural health experts. As a result of that assessment, a generic dashboard view was designed. Its purpose is to summarize key statistics of the envisioned behavioural health interventions for self-monitoring purposes (e.g. steps achieved per day, intervention progress or goals achieved). Based on this generic

mock-up, we implemented a native chat app for Android smartphones for the MobileCoach platform. Figures 1-3 show the graphical user interface of the chat app.




Fig 1. Dashboard view, individual caregiver chat channel PathMate and channel with chatbot Anna



Fig 2. Chatbot Anna, predefined answer options and sensor integration; steps are tracked and used in the chat




Fig 3. Chat channel with the caregiver; the PathMate study team of the children's hospital

3 Significance to Research and Practice

The mobile chat app presented in this paper allows behavioral scientists and health professionals to enrich self-report data with objective sensor data in the everyday life of their clients. This paves the way for a better understanding of whether psychological/self-report data and physiological/objective data are rather complement or alternative measures, a recent research question in the field of NeuroIS [13].

Moreover, it is by far not clear how to design and frame chatbots for DHIs (e.g., as an expert or a “patient like me”) and its interplay with a “physical” caregiver such that they have a positive effect on the bond between caregiver and their clients and thus, also on therapeutic outcomes [6]. In contrast to general purpose chat agents such as Siri (Apple), Alexa (Amazon) or Cortana (Microsoft) and agents with a health focus such as Florence (getflorence.co.uk), Molly (sense.ly) or Lark (lark.com), our chat app allows full control of personal health data and a generic framework to manipulate the design and communication style of chatbots in lab and field settings.

Finally and consistent with the MobileCoach platform, the chat app will be made open source under the Apache 2.0 license to enable a community-driven design such that research teams and (business) organizations interested in chat-based digital coaching approaches do not have to start from scratch but can re-use, revise and improve the existing code together with the MobileCoach platform.

4 Evaluation of the Artifact

Based on prior work demonstrating the acceptance of chat apps by adolescents [11], the first test of the novel chat app was conducted in a children’s hospital in December 2016 with 11 patients ($age_{Mean}=12.6$ years, $SD = 2.4$; 8 girls), who participated in an intervention for the treatment of obesity. The goal of this test was (1) to assess enjoyment, ease of use, usefulness and the intention to use the app [10], and (2) to identify and address major usability problems with the app [12] prior to a randomized controlled trial (RCT), in which the efficacy of a chat-based six-month DHI for the treatment of childhood obesity will be compared to a control group.

First, a chat-based DHI was collaboratively designed by computer scientists, physicians, a psychotherapist, diet and sport experts. The patients were then asked to select a chatbot of their liking, i.e. they could choose between a female and male chatbot (Anna or Lukas). Then, they interacted with the bot for 10 minutes including various chat-based photo, physical activity and quiz interactions. The patients were observed during these interactions by a computer scientist and physician. Afterwards, patients were asked to fill out a questionnaire to assess the app and to provide qualitative feedback on their experience with the app. Similar to prior work [10], we assessed technology perceptions and behavioral intentions with seven-point Likert scales anchored from strongly disagree (1) to strongly agree (7). As young patients deserve special consideration, we used single-item measures to reduce the burden of evaluation [3].

The descriptive statistics of the evaluation are shown in Table 1. Results indicate that the chat app was perceived positive regarding all four constructs. A sign test against the neutral Likert-scale median of 4 supports this observation. Finally, we found no major usability problems based on the observations and the qualitative feedback.

Table 1. Descriptive statistics and results of a sign test against the neutral value 4 on a 7-point Likert-scale ($N=11$). Note: Perceived ease of use (PEU), Perceived enjoyment (PEN), Perceived usefulness (PU) and Intention to use (IU); Significance * / ** / *** $p < .05 / .01 / .001$

#	Item	Mean	Median	SD	p-value
PEU	I found the chat easy to use.	6.7	7.0	0.7	***
PEN	I enjoyed chatting.	6.2	7.0	1.5	**
PU	Chatting with Lukas/Anna could motivate me to accomplish my intervention tasks.	5.9	6.0	1.1	*
IU	I could imagine chatting daily that way.	5.6	6.0	1.4	*

First findings of the aforementioned RCT show that new young patients assigned to the chat-based DHI ($N=14$) completed successfully more than 60% of the daily intervention tasks over the first two months. The efficacy of this DHI will be finally measured by the Body Mass Index after the six-month RCT. We hypothesize that the chat-based DHI is more effective as the chatbot can provide everyday support on therapy goals and tasks, thus increasing therapy adherence compared to patients of the treatment-as-usual control group without everyday support.

In our future work, we will test chat-based DHIs with older patient populations and different therapies to assess the degree to which our findings can be generalized.

Acknowledgements: We would like to thank the CSS Insurance and the Swiss National Science Foundation for their support through grants 159289 and 162724.

References

1. Agarwal, R., Gao, G., DesRoches, C., et al.: The Digital Transformation of Healthcare: Current Status and the Road Ahead. *Information Systems Research* 21, 796-809 (2010).
2. Aluttis, C., Bishaw, T., Frank, M.W.: The workforce for health in a globalized context – global shortages and international migration. *Global Health Action* 7, (2014).
3. Bergkvist, L., Rossiter, J.R.: The Predictive Validity of Multiple-Item versus Single-Item Measures of the Same Constructs. *Journal of Marketing Research* 44, 175-184 (2007).
4. Chaffey, D.: *Mobile Marketing Statistics* compilation, <http://www.smartinsights.com/>, accessed on Feb 2017 (2016).
5. Filler, A., Kowatsch, T., Haug, S., et al.: MobileCoach: A Novel Open Source Platform for the Design of Evidence-based, Scalable and Low-Cost Behavioral Health Interventions. 14th Annual Wireless Telecommunications Symposium (WTS 2015), New York (2015).
6. Flückiger, C., Del Re, A.C., Wampold, B.E., et al.: How central is the alliance in psychotherapy? A multilevel longitudinal meta-analysis. *Journal of Counseling Psychology* 59, 10-17 (2012).
7. Haug, S., Kowatsch, T., Paz Castro, R., et al.: Efficacy of a web- and text messaging-based intervention to reduce problem drinking in young people: study protocol of a cluster-randomised controlled trial. *BMC Public Health* 14, 1-8 (2014).
8. Haug, S., Paz Castro, R., Filler, A., et al.: Efficacy of an internet and SMS-based integrated smoking cessation and alcohol intervention for smoking cessation in young people: study protocol of a two-arm cluster randomised controlled trial. *BMC Public Health* 14, (2014).
9. Haug, S., Paz, R., Kowatsch, T., et al.: Efficacy of a web- and text messaging-based intervention to reduce problem drinking in adolescents: Results of a cluster-randomised controlled trial. *Journal of Consulting and Clinical Psychology* 85, 147-159 (2017).
10. Kamis, A., Koufaris, M., Stern, T.: Using an Attribute-Based Decision Support System for User-Customized Products Online: An Experimental Investigation. *MIS Quarterly* 32, 159-177 (2008).
11. Lilian, S., Gregor, W., Sarah, G., et al.: MIKE - Medien, Interaktion, Kinder, Eltern. Zürcher Hochschule für Angewandte Wissenschaften, Zürich (2015).
12. Nielsen, J., Landauer, T.K.: A mathematical model of the finding of usability problems. CHI '93 Proceedings of the INTERACT '93 and CHI '93 Conference on Human Factors in Computing Systems, pp. 206-213. ACM, Amsterdam, The Netherlands (1993).
13. Tams, S., Hill, K., de Guinea, A.O., et al.: NeuroIS - Alternative or Complement to Existing Methods? Illustrating the Holistic Effects of Neuroscience and Self-Reported Data in the Context of Technostress Research. *Journal of the Association for Information Systems* 15, Article 1 (2014).
14. WHO: Global Health and Aging. World Health Organization, Genève, Switzerland (2011).
15. WHO: Draft comprehensive global monitoring framework and targets for the prevention and control of noncommunicable diseases. World Health Organization, Genève, Switzerland (2013).